

CLENIUM

Composition:
Each uncoated tablet contains:
Clenbuterol HCL E.P. 40mcg
Excipients q.s.
Clenbuterol is a sympathomimetic amine used by sufferers of breathing disorders as a decongestant and bronchodilator. People with chron breathing disorders such as asthma use this as a bronchodilator to make breathing easier. Clenbuterol is a B2 agonist with some structural and pharmacological similarities to epinphrine and salbutamol, but its effects are more potent and longer lasting as a stimulant and thermogenic drug. It causes an increase in aerobic capacity, central nervous system stimulation, blood pressure, and oxygen transportation. It increases the rate at which body fat is metabolized while increasing the body's basal metabolic rate (BMR). It is commonly used for smooth muscle relaxant properties as a bronchodilator and tocolytic.



PLAZOL

Composition:
Each uncoated tablet contains:
Stanozolole U.S.P 10mg
Excipients q.s.
Stanozol is al 7-alpha-alkylated anabolic steroid with good oral bioavailability While stanozol's androgenic activity Index is nearly the same as testosterone, its myotrophic activity is much higher at 2.0-3.7. Stanozol does not aromatize and does not cause water retention. Stanozol has been used for weight loss management in HIV patients with chronic obstructive pulmonary disease and has showed significant improvement in weight, body mass index, LBM, and muscle size. Stanozol has been shown in vitro to significantly enhance collagen synthesis,A positive effect of stanazolol on wound healing in patients with non-healing wounds has also been demonstrated. Clinical studies show that the treatment with stanozol during the induction phase of chemotherapy results in a positive effect on the duration of cancer free remission.



ANAVAR

Composition:
Each uncoated tablet contains:
Oxandrolone U.S.P 10mg
Excipients q.s.
Oxandroloneis classified as a Class I anabolic steroid, which when combined with Class II compounds such as Dianabol or Anadrol becomes more effective. It does not contribute much with the use of high doses of class I steroids such as trenbolone or testosterone considered as a steroid of mixed activity. However, its use with moderate doses of testosterone can be useful, although it is an expensive option.Anavar is often considered a weak steroid.This is partly because when only a Class I steroid is used, its effectiveness is never total. Another cause is that bodybuilders as well as experts in the field sometimes make unfortunate and unfair comparisons when dealing with anabolic steroids. If, for example, 8 tablets a day give a meager result, then it is thought that the drug is weak or does not work. In addition, oxandrolone has been available as Anavar at 2.5 mg which only demonstrates a daily dosage of 20 mg or the equivalent of 140 mg per week.



DYBOL

Composition:
Each uncoated tablet contains:
Methandienone IH 10mg
Excipients q.s.
Methandrostenolone is a 17-alpha alkylated oral anabolic-androgenic steroid with anabolic properties that are more pronounced than its androgenic effects. It has little progestational activity and easily converts to estrogens, Methandrostenolone has an anabolic activity index 0.60 and an androgenic activity index 0.20.Methandrostenolone has an extra double bond in the A-ring and, therefore, is not converted by 5-alpha reduction into DHT.Since methandrostenolone was created in the 1960s, it has been used for many different applications. At present, the only legitimate therapeutic indications for this anabolic steroid are: replacement of male sex steroids in men who have androgen deficiency, for example as a result of loss of both testes: the treatment of certain rare forms of aplastic anemia which are or may be responsive to anabolic androgens: and, In certain countries, to counteract catabolic states, for example after major trauma.



HALOTESTIN

Composition:
Each uncoated tablet contains:
Fluoxymesterone U.S.P 5mg
Excipients q.s.
INDICATIONS:
Certain cases of disseminated breast cancer in women.
Osteoporosis due to androgen deficiency in hypogonadal males.
CONTRA-INDICATIONS:
Not intended for use in children.
Known or suspected prostatic carcinoma and mammary carcinoma in the male.
Not intended for use in female patients other than those with disseminated breast cancer.
Contraindicated in nephrosis or the nephrotic phase of nephritis, cardiac and renal failure, hypercalcaemia, oedema, jaundice, liver disease with impaired bilirubin excretion, testicular and hepatic carcinoma.
DOSAGE AND DIRECTIONS FOR USE:
Muscle anabelism: 10 - 30 mg taken orally each day in divided doses for duration of 4-6 weeks. Hereditary angioedema: The dosage requirement for continuous treatment of hereditary angioedema with Flubolic of the clinical response of the patient. Commonly 2-6 mg taken orally each day should be individe in divided doses initially and reduced by physician thereafter.
Effective dose for male: 5 - 20 mg per day
Effective dose for female: 10-40 mg per day
OTHER ADVERSE REACTIONS MAY INCLUDE:
Oligospermin and decreased ejaculatory volume;
Suppression of ovarian activity, atrophy of the breasts and endometrial tissue.
Amenorrhoea and inhibition of spermatogenesis.
Water and salt retention.
Premature epiphyseal closure.
If signs of virilisation develop, treatment should be discontinued.
Increase in nitrogen retention and skeletal weight;
PATIENTS WITH THE FOLLOWING CONDITIONS SHOULD BE MONITORED:
Latent or overt cardiac failure, renal dysfunction, hypertension, epilepsy or migraine (or a history of these conditions), since anabolic steroids may induce salt and fluid retention; diabetes, since anabolic steroids may improve the glucose tolerance and decrease the need for insulin or other antidiabetic drugs; incomplete statural growth, since anabolic steroids in high dosages may accelerate epiphyseal closure; skeletal metastases, since anabolic steroids may induce hypercalcaemia and hypercalciuria in these patients. liver dysfunction.



PROVIRON

Composition:
Each uncoated tablet contains:
Mesterololn U.S.P 25mg
Excipients q.s.
Before you start this treatment, read the manufacturer's printed information leaflet from inside your pack. The leaflet will give you more information about mesterolone tablets.
Take the tablets exactly as your doctor has told you to. To begin with, it is usual to be prescribed 25 mg (one tablet), to be taken three or four times a day. If you need to continue taking the tablets for more than a few months, your doctor may reduce the number of tablets you need to take each day. Your doctor will tell you what dose is right for you and this will also be printed on the label of the pack to remind you.
Swallow the tablet whole with a drink of water. You can take mesterolone before or after meals.
Try to take your doses at the same times each day, as this will help you to remember to take them.
If you forget to take a dose, you should take it when you remember. However, if when you remember, it is nearly time for your next dose then leave out the forgotten dose. Do not take two doses at the same time to make up for a missed dose.
Important: if you experience frequent erections or erections that last for a long time, you must let your doctor know about this. Your treatment will need to be reviewed in order to prevent any injury.If you experience any other symptoms which you think may be due to this medicine, speak with your doctor or pharmacist.How to store mesteroloneKeep all medicines out of the reach and sight of children.Store in a cool, dry place, away from direct heat and light.

